## Discovery of the "Compton Shoulder" in the Iron Line from GX301–2

#### Shin Watanabe(ISAS)

M. Sako, M. Ishida, Y. Ishisaki, S. Kahn, T. Kohmura. U. Morita,

F. Nagase, F. Paerels, T. Takahashi

Introduction(Compton scattering)

When an X-ray photon ( $E_0$ ) is Compton-scattered by an electron in a cold material,

$$E_1 = \frac{E_0}{1 + E_0 / mc^2 \left( 1 - \cos \theta \right)}$$

If any X-ray emission lines are observed,

"Compton shoulder"

- the direct evidence of Compton scattering
- a probe for the physical states of the materials

Observations with high energy resolution hold the key!

Iron K $\alpha$  line(6.4keV) -> Compton shoulder: 6.24-6.40keV


width: 160eV

Wave length

Energy

Chandra HETG is capable of detecting Compton shoulders

## Observation of GX301–2 with Chandra HETG




GX301-2

- High Mass X-ray Binary (NS and super-giant(B2lae))
- Strong iron Kα line (EW ~ 1keV)
   -> the most promising target!!

We have observed at 3 orbital phases (Intermediate(0.167-0.179) ~40ksec
Near-Apastron(0.480-0.497) ~60ksec
Pre-Periastron(0.970-0.982) ~40ksec)

#### Result of the observation


Energy spectrum from HEG ±1 order data



A "clear" Compton shoulder is detected for the first time!

## Variability of the Compton shoulder

Light curve of HEG ±1 order



The first ~20 ksec The second ~20 ksec

### How to reproduce the shoulder

#### Issues to be considered

- Two competing physical processes (photoelectric absorption and Compton scattering)
- Multiple scatterings
   N<sub>H</sub> ~ 10<sup>24</sup>cm<sup>-2</sup> —> Optical depth of each process is ~1
- Geometry




It is too complex for analytical approaches.

The easiest way is the Monte Carlo simulation of photon transport.

#### The Monte Carlo simulation

- A homogeneous spherical cloud
  - •The NS (the power-law X-ray source) is located at the center


Photoelectric absorption and Compton scattering




## Results of Monte Carlo simulations(1)

A transition when NH is changed. (kTe is fixed 0eV)

(Metal Abundance: 0.7 cosmic)





The flux ratio of the shoulder to the line vs NH


Metal abundance for cosmic abundance (Feldman 1992)

0.5 x cosmic 0.7 x cosmic 1.0 x cosmic

## Results of Monte Carlo simulations(2)

A transition when kTe is changed. (NH is fixed 1x10<sup>24</sup> cm<sup>-2</sup>)


(Metal Abundance: 0.7 cosmic)



#### The flux ratio of the shoulder to the line

- → the optical depth and the abundance of the cloud
  The shape of the shoulder
  - the temperature of scattering electrons

## Fitting of the Compton shoulder




0.7 cosmic abundance is adopted as the metal abundance

 $N_{H}$  11.7+2.3/-2.0 x10<sup>23</sup> cm<sup>-2</sup>, 8.4±1.6 x10<sup>23</sup> cm<sup>-2</sup> Upper limit of kTe < 3.7eV, <0.7eV (90%)

The new means for probing materials around X-ray sources!

# N<sub>H</sub> from the Compton shoulder can reproduce the whole spectrum shape



Full Monte Carlo simulation

- a spherical cloud
- a power-law source with  $\Gamma$ =1.0
- a metal abundance: 0.7 cosmic
- N<sub>H</sub> derived from the Compton shoulder

We can obtain a self-consistent solution from the analysis of the Compton shoulder!

## Summary

- We discovered the Compton shoulder in the iron Kα line of GX301–2
- From properties of the Compton shoulder, we have obtained physical parameters characterizing the material around the X-ray source, using Monte Carlo simulations.
- The new means for probing cold materials using Compton shoulders was established.

#### **Future**

#### **ASTRO-E2 XRS**

- ◆ Larger effective area(300cm<sup>2</sup>@6keV)
  - -> other X-ray binaries, AGNs diffuse sources (the Galactic center region)
- Excellent energy resolution (6eV@6keV)
  - -> geometrical conditions